

Neural, electrophysiological and anatomical basis of brain-network variability and its characteristic changes in mental disorders

Jie Zhang,^{1,2,*} Wei Cheng,^{1,*} Zhaowen Liu,^{1,3,*} Kai Zhang,^{4,*} Xu Lei,^{5,6} Ye Yao,¹ Benjamin Becker,⁷ Yicen Liu,¹ Keith M. Kendrick,⁷ Guangming Lu^{2,8} and Jianfeng Feng^{1,9,10,11}

*These authors contributed equally to this work.

See Mattar *et al.* (doi:10.1093/aww151) for a scientific commentary on this article.

Functional brain networks demonstrate significant temporal variability and dynamic reconfiguration even in the resting state. Currently, most studies investigate temporal variability of brain networks at the scale of single (micro) or whole-brain (macro) connectivity. However, the mechanism underlying time-varying properties remains unclear, as the coupling between brain network variability and neural activity is not readily apparent when analysed at either micro or macroscales. We propose an intermediate (meso) scale analysis and characterize temporal variability of the functional architecture associated with a particular region. This yields a topography of variability that reflects the whole-brain and, most importantly, creates an analytical framework to establish the fundamental relationship between variability of regional functional architecture and its neural activity or structural connectivity. We find that temporal variability reflects the dynamical reconfiguration of a brain region into distinct functional modules at different times and may be indicative of brain flexibility and adaptability. Primary and unimodal sensory-motor cortices demonstrate low temporal variability, while transmodal areas, including heteromodal association areas and limbic system, demonstrate the high variability. In particular, regions with highest variability such as hippocampus/parahippocampus, inferior and middle temporal gyrus, olfactory gyrus and caudate are all related to learning, suggesting that the temporal variability may indicate the level of brain adaptability. With simultaneously recorded electroencephalography/functional magnetic resonance imaging and functional magnetic resonance imaging/diffusion tensor imaging data, we also find that variability of regional functional architecture is modulated by local blood oxygen level-dependent activity and α -band oscillation, and is governed by the ratio of intra- to inter-community structural connectivity. Application of the mesoscale variability measure to multicentre datasets of three mental disorders and matched controls involving 1180 subjects reveals that those regions demonstrating extreme, i.e. highest/lowest variability in controls are most liable to change in mental disorders. Specifically, we draw attention to the identification of diametrically opposing patterns of variability changes between schizophrenia and attention deficit hyperactivity disorder/autism. Regions of the default-mode network demonstrate lower variability in patients with schizophrenia, but high variability in patients with autism/attention deficit hyperactivity disorder, compared with respective controls. In contrast, subcortical regions, especially the thalamus, show higher variability in schizophrenia patients, but lower variability in patients with attention deficit hyperactivity disorder. The changes in variability of these regions are also closely related to symptom scores. Our work provides insights into the dynamic organization of the resting brain and how it changes in brain disorders. The nodal variability measure may also be potentially useful as a predictor for learning and neural rehabilitation.

1 Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai 200433, PR China

2 Department of Medical Imaging, Jinling Hospital, Nanjing University School of Medicine, Nanjing 210002, PR China

Received January 7, 2016. Revised March 18, 2016. Accepted April 26, 2016.

© The Author (2016). Published by Oxford University Press on behalf of the Guarantors of Brain. All rights reserved.

For Permissions, please email: journals.permissions@oup.com

- 3 School of Computer Science and Technology, Xidian University, Xi'an 710071, Shanxi, PR China
- 4 NEC Laboratories America, 4 Independence Way, Princeton, NJ 08540, USA
- 5 Key Laboratory of Cognition and Personality, Ministry of Education, Chongqing, China
- 6 Sleep and NeuroImaging Center, Faculty of Psychology, Southwest University, Chongqing, China
- 7 Key Laboratory for Neuroinformatics, Center for Information in Medicine, University of Electronic Science and Technology of China, Chengdu, PR China
- 8 Jinling Hospital-Fudan University Computational Translational Medicine Center, Jinling Hospital, Nanjing University School of Medicine, Nanjing 210002, PR China
- 9 Department of Computer Science, University of Warwick, Coventry CV4 7AL, UK
- 10 Collaborative Innovation Center for Brain Science, Fudan University, Shanghai, 200433, PR China
- 11 Shanghai Center for Mathematical Sciences, Shanghai, 200433, P.R. China

Correspondence to: Professor Jianfeng Feng,
 Department of Computer Science,
 University of Warwick,
 Coventry CV4 7AL,
 UK
 E-mail: jianfeng64@gmail.com

Correspondence may also be addressed to: Prof. Jie Zhang, Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai 200433, China E-mail: jzhang080@gmail.com

Keywords: resting-state functional MRI; functional brain networks; temporal variability; brain flexibility and adaptability; mental disorders

Abbreviations: ADHD = attention deficit hyperactivity disorder; BOLD = blood oxygen level-dependent; DMN = default-mode network; DTI = diffusion tensor imaging; RIIC = ratio of intra- to intercommunity structural connection

Introduction

The human brain demonstrates remarkable variability in its structure and function (Rademacher *et al.*, 2001; Sugiura *et al.*, 2007; Frost and Goebel, 2012), which explains inter-subject variability in cognitive function and other behaviours. Intersubject variability in resting state functional connectivity is heterogeneous across the cortex, and it is significantly correlated with the degree of evolutionary cortical expansion (Mueller *et al.*, 2013). Individual variability in functional connectivity is also predictive of task performance (Baldassarre *et al.*, 2012). Recently, the temporal variability of neuronal activity and functional connectivity/functional networks has attracted increasing attention (Bassett *et al.*, 2011, 2013, 2015; Hutchison *et al.*, 2013a; Mueller *et al.*, 2013; Calhoun *et al.*, 2014; Kopell *et al.*, 2014; Kucyi and Davis, 2014; Tagliazucchi and Laufs, 2014; Braun *et al.*, 2015). For example, variability of blood oxygen level-dependent (BOLD) signal, which was previously considered to be measurement-related 'noise', has been demonstrated to have significant age-predictive power (Garrett *et al.*, 2010) and is related to task performance (Garrett *et al.*, 2011). Even during rest, either sliding window or time-frequency analysis shows non-stationarity in both spontaneous brain activity (McIntosh *et al.*, 2008; Lippe *et al.*, 2009; Garrett *et al.*, 2010, 2011; Misic *et al.*, 2010; Protzner *et al.*, 2010; Samanez-Larkin *et al.*, 2010) and interactions among brain regions (Chang and Glover, 2010; Kang *et al.*, 2011; Majeed *et al.*, 2011; Hutchison *et al.*, 2013b; Mueller *et al.*, 2013).

To date, most work on dynamic brain network analysis either focuses on single functional connectivity between a given pair of regions of interest (Chang and Glover, 2010; Kang *et al.*, 2011; Majeed *et al.*, 2011; Hutchison *et al.*, 2013a; Kucyi and Davis, 2014; Zalesky *et al.*, 2014), or the connectivity of the whole brain (Allen *et al.*, 2014). Temporal variability of brain networks at the mesoscale, i.e. the functional architecture of a given region (defined as the overall functional connectivity profile associated with the region), has never been investigated. While non-stationarity in functional connectivity/networks has been revealed, the underlying mechanisms and neuroanatomical basis for temporal variability are still unknown. The advantage of investigating temporal variability of functional architecture associated with a specific brain region (i.e. a mesoscale analysis), is 2-fold. First, it allows coupling between temporal variability of the functional architecture of a region and its neural activity to be conveniently analysed. This helps delineate factors contributing to temporal variability, thus shedding light on the underlying mechanisms. Second, it facilitates the construction of a whole-brain topography of variability, which allows localization of regions showing significant variability changes in patients, thus helping to define the dynamics of functional brain networks for various brain disorders. In comparison, analysis at the level of single functional connectivity would increase the burden of correction for multiple comparisons, while analysis of the whole-brain functional connectivity simultaneously may be less sensitive in uncovering local (regional) changes.

Temporal variability of regional, or nodal functional architecture can be characterized by first constructing whole-brain functional networks from BOLD signals at successive, non-overlapping time windows and second, by comparing the functional architecture of a region of interest across different windows. Accordingly, we hypothesized that temporal variability of regional functional architecture is modulated by local neural activity, as manifested by the BOLD signal and EEG, and that it has an anatomical substrate. Using simultaneously recorded EEG/functional MRI and functional MRI/diffusion tensor imaging (DTI) data, we found that the temporal variability of functional architecture of a given region is modulated by local BOLD activity, i.e. amplitude and frequency of BOLD oscillations during the scan, and that it is positively associated with the EEG α power of the region of interest. Variability is also related to the ratio of intra- to inter-community structural connectivity of a region. Application to different psychiatric disorders, including schizophrenia, autism spectrum disorder, and attention deficit hyperactivity disorder (ADHD) with their matched controls, revealed disease-specific variability changes in the default mode network (DMN), as well as visual and subcortical regions of the brain, which provides new insights of spatiotemporal organization of the brain networks and how it changes in patients with psychiatric disease.

Materials and methods

Participants, image acquisition and data preprocessing

The study included eight multicentre datasets involving 1180 subjects. Six resting state functional MRI datasets were used for case-control studies of variability change in patients with psychiatric disorders: schizophrenia [Dataset 1: Taiwan (Guo *et al.*, 2014); Dataset 2: COBRE], autism (Dataset 3: New York University-NYU; and Dataset 4: University of Melbourne-UM, which are from ABIDE Consortium http://fcon_1000.projects.nitrc.org/indi/abide/) and ADHD (Dataset 5: Peking University-PKU; and Dataset 6: New York University-NYU, which are part of the 1000 Functional Connectome Project http://fcon_1000.projects.nitrc.org/indi/adhd200/) and matched controls. Demographic details and medication information are given in Table 1 and Supplementary Table 1. An EEG/functional MRI dataset (Dataset 7) and a functional MRI/DTI dataset (Dataset 8) from IMAGEN consortium (Schumann *et al.*, 2010) were also used to investigate the electrophysiological and structural basis of variability, respectively. Details of participants, image acquisition and data preprocessing can be found in the Supplementary material, which also includes a discussion of global signal removal and details of data scrubbing.

In view of the fact that we combined multicentre data for case-control studies, and this might possibly result in a large variation, we set up the following exclusion criteria to help ensure data quality: (i) subjects with poor structural scans, or functional MRI data, making successful preprocessing

unlikely [i.e. normalization to Montreal Neurological Institute (MNI) space], or without complete demographic information; and (ii) head movement, including subjects with >10% displaced frames in a scrubbing procedure (Power *et al.*, 2014; Cheng *et al.*, 2015b), or maximal motion between volumes in each direction >3 mm, and rotation about each axis >3° (Cheng *et al.*, 2015a). In each dataset, patients and controls were screened so that the total root mean square displacements did not show significant differences. For the ADHD dataset, we only have preprocessed data (i.e. BOLD time series) from the public website, thus the data scrubbing procedure cannot be performed. For the autism dataset, subjects with an overall IQ score exceeding 2 standard deviations (SD) from the overall ABIDE sample mean were not included.

Temporal variability of regional functional architecture

To characterize the temporal variability of the functional architecture associated with a given region (Fig. 1A), we first segmented all BOLD time series into n non-overlapping windows each with length l . Within the i th time window, the whole-brain functional network F_i (an $m \times m$ matrix, with m nodes) is obtained using Pearson correlation as the measure of functional connectivity. The functional architecture of a region k at time window i is defined as the overall functional-connectivity profile of region k , i.e. $F_i(k, :)$ which is a m -dimensional vector and is shortened as $F_{i,k}$. We then define the variability of a region of interest k as:

$$V_k = 1 - \overline{\text{corrcoef}(F_{i,k}, F_{j,k})}, i, j = 1, 2, 3, \dots, n, i \neq j, \quad (1)$$

as illustrated in Fig. 1A. The latter part of V_k compares the functional architecture, i.e. overall functional connectivity profile associated with brain region k across different time windows, which is the averaged correlation coefficient among different functional architecture of region k and thus a similarity measure. Next, a deduction from 1 indicates temporal variability of a region. In this way, it is possible to both evaluate temporal variability of the functional architecture at the network level and simultaneously localize this to a specific brain region. In fact, this approach has been used in a different context to measure intersubject variability of regional functional architecture (Mueller *et al.*, 2013). We have adopted this same measure to address the temporal variability of the functional architecture of a region in one subject.

To reduce the influence from segmentation scheme for the BOLD signal, we perform $l - 1$ times segmentation for a given window length l , and each time we start from the s th point ($s = 1, 2, \dots, l$) and average the variability obtained from $l - 1$ times segmentation. To avoid arbitrary choice of window length in the applications, we calculate V_k at different l ($l = 10, 11, 12, \dots, 20$ volumes, equal to 20, 22, 24, ... 40 s) and then take the average value as the final variability of the region of interest. We chose the above window length as it was suggested that window sizes around 30–60 s produce robust results in image acquisitions, cognitive states (Shirer *et al.*, 2012) and topological descriptions of brain networks (Jones *et al.*, 2012). In fact, we found that variability obtained at different window lengths (e.g. 20 s, 30 s, 40 s) was highly correlated ($r > 0.98$, Supplementary Fig. 1), indicating that this metric

Table 1 Demographic information for the eight datasets used, involving three mental disorders and two multimodal datasets from healthy controls

		Groups	n	Age, years	Sex (M/F)	PANSS (P)	PANSS (N)	PANSS (G)	Illness duration
SCZ	Taiwan dataset (#1)	Controls	62	29.9 ± 8.6	25/37				
		Schizophrenia	69	31.9 ± 9.6	35/34	11.9 ± 4.7	13.6 ± 6.3	27.3 ± 9.6	7.2 ± 6.6
	COBRE dataset (#2)	Controls	67	34.8 ± 11.3	42/11				
		Schizophrenia	53	36.8 ± 13.7	46/21	14.9 ± 4.6	14.7 ± 5.2	29.7 ± 8.2	8.9 ± 6.9
Autism	NYU dataset (#3)	Controls	102	15.9 ± 6.	76/26	21.4 ± 12.7			
		Autism	75	14.8 ± 7.0	65/10	92.6 ± 31.0			
	UM dataset (#4)	Controls	64	15.1 ± 3.7	48/16				
		Autism	38	13.6 ± 2.4	31/7				
ADHD	PKU dataset (#5)	Control	143	11.4 ± 1.9	84/59	29.3 ± 6.4			
		ADHD	99	12.1 ± 2.0	89/10	50.4 ± 8.2			
	NYU dataset (#6)	Control	108	12.2 ± 3.1	54/54	45.4 ± 6.0			
		ADHD	140	11.1 ± 2.7	106/34	71.9 ± 8.7			
Multi-modal imaging	EEG/functional MRI (#7)	Control	26	21.4 ± 2.0	15/11				
	Functional MRI/DTI (IMAGEN, #8)	Control	142	14.5 ± 0.2	66/76				

ADOS = Autism Diagnostic Observation Schedule; SCZ = schizophrenia.




Figure 1 Flow chart of temporal variability analysis. (A) Definition of variability for a given region k . The time series are BOLD signals for all brain regions, which are segmented into non-overlapping windows. Functional brain networks are constructed in each time window. The variability of region k is obtained by comparing functional connectivity profile of region k at different windows. (B) Correlation analysis of temporal variability with measures from multimodal neuroimaging data, such as simultaneously recorded EEG and DTI data. (C) Case-control studies for different mental disorders, including schizophrenia, autism and ADHD using temporal variability.

does not necessarily depend on the choice of window length, and we take the average variability over different window lengths as mentioned above.

Correlation analysis between temporal variability and local/global measures from multimodalities

To identify the neural, electrophysiological and anatomical basis of temporal variability, we performed extensive correlation analyses using BOLD, EEG/functional MRI and functional MRI/DTI data collected from healthy controls (Fig. 1B). To establish the neural basis, the variability of a region was correlated with its BOLD activity (the variance of BOLD signal during the whole scan) and node degree across 90 brain regions for each subject in Dataset 5 (which has the largest number of controls). To establish the electrophysiological basis of variability, we performed correlation analyses between the variability of a region and its α band power of simultaneously recorded EEG during the entire scan (in Dataset 7, across all 28 electrodes, see [Supplementary material](#) for details of the 28 corresponding regions in the AAL template). To establish the anatomical basis, we performed correlation analyses between the variability of a region and the ratio of intra- to intercommunity structural connection (RIIC) across 90 brain regions for each subject in the IMAGEN dataset (Dataset 8). A high RIIC implies that the region connects more with regions belonging to the same functional module, while a low RIIC means the region connects more with nodes belonging to communities other than its own (see DTI data preprocessing in the [Supplementary material](#) for details).

Case-control studies: meta-analysis integration of multicentre results

To identify regions with significant change in variability for various mental disorders including schizophrenia, autism and ADHD, we first performed *t*-test between patients and matched healthy controls for all 90 brain regions (AAL template) for each dataset, with age, sex and root mean square displacements of head movement being regressed out (Fig. 1C). For disorders with multiple datasets, we then used the Liptak-Stouffer *z*-score method (Liptak, 1958) to integrate the results (e.g. MRI; Glahn *et al.*, 2008): the *P*-value of each region in the relevant dataset *i* was converted to the corresponding *z* score: $z_i = \Phi^{-1}(1 - p_i)$, where Φ is the standard normal cumulative distribution function. Then a combined *z*-score for a functional connectivity was obtained using the Liptak-Stouffer formula as:

$$Z = \frac{\sum_{i=1}^k w_i z_i}{\sqrt{\sum_{i=1}^k w_i^2}}, \quad (2)$$

where w_i is the inverse of the variance of z_i . *Z* follows a standard normal distribution under the null hypothesis and is transformed into its corresponding *P*-value with both Bonferroni ($P = 0.05$) and false discovery rate (FDR) ($q = 0.05$) correction used to correct for multiple comparisons.

Results

Variability reflects change of community membership, or flexibility of a region

The temporal variability of a region defined in our paper characterizes the collective changes of all its functional connectivities over time. Low temporal variability means that this functional architecture of a given region of interest is highly correlated across different time windows, or alternatively that the dynamical functional connectivity time series between the region of interest and all other regions are highly synchronized (Fig. 3A and [Supplementary Fig. 2A](#)). On the contrary, high temporal variability means that the dynamical functional connectivity series between a region of interest and other regions remains independent (Fig. 3A and [Supplementary Fig. 2B](#)). Due to the dynamic nature of the functional brain network in resting state, a region of interest may connect with different brain regions and be involved in different functional communities/modules at different times. The variability herein defined is also a good indicator for this property ([Supplementary Fig. 3](#)). Of the 62 healthy control subjects in Dataset 1, all subjects showed significant negative correlation between the variability of a region and stability of its intra-community members ($P < 0.05$). The intra-community members of a region of interest indicate the regions belonging to the same functional module with the region of interest. The above negative correlation suggests that the higher the variability of a region of interest the less stable its intra-community members, i.e. its intra-community members change frequently with time. Since the intra-community members of a region of interest are not stable, the region of interest may connect with different brain regions, and functional communities at different times. This result indicates that the temporal variability we defined reflects the ability or tendency of a region to reconfigure itself into different functional communities, or its flexibility in terms of functional integration/coordination with different neural systems. The larger the temporal variability of a region of interest, the more functional communities it will be involved with at different times (see [Supplementary Fig. 3](#) for details).

Stable brain-wide topography of variability in healthy control subjects

For healthy control subjects, we found a non-uniform distribution of variability throughout the brain (Fig. 2 and [Supplementary Table 2](#)). Various datasets consistently demonstrate low variability in primary sensory cortices (e.g. Heschl's gyri, postcentral and calcarine gyrus), visceral sensory cortex (insular), unimodal association cortex (middle/superior occipital gyrus, cuneus, lingual gyrus, superior temporal gyrus), and default mode systems such as the




Figure 2 Whole-brain variability topography on AAL template for healthy controls. The variability is averaged over the results obtained from controls in six different datasets (Datasets 1–6). See Supplementary Table 2 for details.

medial frontal gyrus and posterior cingulate/precuneus. In comparison, the transmodal association cortices, including the heteromodal association cortex (such as the anterior association cortex, which includes interior and orbital frontal gyrus, the posterior association cortex including inferior parietal gyrus, inferior/middle temporal gyrus and paracentral lobule) and the limbic association cortex (such as the temporal pole, hippocampus, parahippocampus and amygdala; Pearlson *et al.*, 1996; Bullard *et al.*, 2013) all demonstrate high variability. See Supplementary Table 3 for details of the 90 brain regions in AAL template and their characteristics.

As shown in Supplementary Fig. 4, the temporal variability we defined demonstrates great similarity by showing a consistent pattern in healthy controls across Datasets 1–6, indicating the robustness of the variability defined in our work.

Temporal variability as an index of regional adaptability in the brain

The human brain is perhaps the most adaptable and changing part of the body, which accounts for its incredible learning capability. Interestingly, the transmodal cortices of the brain, which are found to demonstrate the highest levels of temporal variability (the top regions in Supplementary Table 2), have almost all been implicated in key aspects of learning, suggesting that it may provide an index of the adaptability and plasticity of these brain regions, which support learning. For example, the hippocampus is extensively involved in many aspects of learning

and memory (Deng *et al.*, 2010), the inferior temporal cortex in visual association learning (Kawasaki and Sheinberg, 2008), the olfactory cortex in olfactory learning (Fletcher and Chen, 2010), and the caudate in reinforcement-based associative learning (Williams and Eskandar, 2006) and classification learning (Seger and Cincotta, 2005). In particular, there is solid evidence that learning-associated neurogenesis (birth of brain cells) can occur in a number of these regions in the adult mammalian brain (Gould *et al.*, 1999; Rakic, 2002), including hippocampus, olfactory bulb and inferior temporal cortex. Our proposed link between high temporal regional variability and plasticity underpinning learning is further supported by our additional finding that it is positively associated with various kinds of IQ score in two independent datasets (see Supplementary Fig. 7 for details).

Correlation between variability and measures from multiple neuro-modalities

Blood oxygenation level-dependent activity

Variability of a region results from the temporal changes of its functional architecture triggered by changes in BOLD signals. We first analysed the correlation between variability and other measures derived from the BOLD signal over the entire period of scan, including local measures such as the amplitude and frequency of BOLD activity, and global measures such as the degree of the region (defined as the number of a region's functional connectivity




Figure 3 Correlation between temporal variability and local/global metrics obtained from various neuroimaging modalities. Green indicates negative correlation among the variables, and red for positive correlation, with the strength of correlation listed in Supplementary Table 4. (A) Temporal variability of the functional architecture of a region of interest. The upper part indicates that the region of interest has high variability and that the multiple dynamical functional-connectivity time-series between the region of interest and the other regions are independent. The lower part indicates that the region of interest has low variability and that its dynamic functional connectivity time series are synchronous. (B) Amplitude of BOLD activity of a region of interest. The upper part is for a region with low amplitude, and lower part is for a region with high amplitude. Here the variance of BOLD signal during the entire period of scan is used to represent the amplitude of BOLD oscillation. (C) α band power of EEG of a region of interest. The upper part shows a region with large α band power, and the lower part shows a region with small α band power. α band power of the EEG of the entire period of scan is used here. (D) Node degree of a region of interest obtained from a BOLD-constructed brain network (defined as the number of functional connectivities of a region with absolute strength larger than 0.3). The upper part shows a region with a low degree and the lower part shows a region with a high degree. (E) Ratio of intra-community to inter-community structural connectivity (RIIC). The upper part shows a region with a small RIIC (i.e. the region of interest connects more with nodes belonging to different communities), and the lower part is for a region with a high RIIC (i.e. the region of interest connects with nodes belonging to the same communities). All upper parts (or lower parts) in each panel are correlated such that a region with low variability will have higher BOLD activity, a high node degree, low α band power and a high RIIC.

whose strength is larger than 0.3). We found that the temporal variability of a region correlates negatively with both the amplitude of its BOLD activity and the node degree, since the BOLD activity of a region and its degree are positively correlated, see (Fig. 3A, B and D). In addition, the variability of a region is negatively associated with the energy of low frequency components of the BOLD signal. For scatter plots and details of correlations, (see Supplementary Fig. 5D, E and F and Supplementary Table 4).

Electrophysiological recording

We next analysed the correlation between the variability of a region and its α band power derived from EEG recorded

simultaneously with functional MRI data. Correlations were performed across brain regions having both BOLD signal and EEG recordings in each of the subjects in Dataset 7. We found that the variability of a region was mostly positively correlated with its α band power during the entire period of scan, with 8 of 26 subjects (31%) showing a significant positive correlation (Fig. 3A and C; detailed in Supplementary Table 5).

Structural connectivity: diffusion tensor imaging

Finally, we analysed the correlation between the variability of a region and its structural connectivity derived from diffusion tensor imaging. We used the RIIC to determine

the extent to which a node is structurally connected to nodes within its own community. Of 142 healthy subjects in the IMAGEN dataset (Dataset 8), 49 showed a significant correlation ($P < 0.05$) between regional variability and regional RIIC, with 40 (28%) showing significant negative correlation ($r = -0.28 \pm 0.0032$; Fig. 3A and E). Therefore, the more intra-community connections a region has (relative to its inter-community connections), the smaller its variability will be. A brain region with more fibre connections to those of the same community would be involved more stably in that functional community, thus showing less variability. In comparison, a brain region structurally connected to regions belonging to many other communities will switch among them, i.e. the region belongs to different functional communities at different times, resulting in high variability.

Disease-specific changes of variability in mental disorders

Next, we performed a whole-brain variability analysis in patients with schizophrenia, autism spectrum disorders (ASD) and ADHD and identified disease-specific changes by comparing them to matched healthy controls. For disorders with more than one dataset, meta-analysis was adopted to integrate results from multiple datasets. For schizophrenia, >20% of brain regions ($n = 19$) showed significant differences in variability (Fig. 4A and Table 2). Variability decreased mainly in DMNs, such as rectus, hippocampus, parahippocampus, inferior parietal gyrus and temporal lobe (middle temporal pole and inferior temporal gyrus), while it increased most prominently in subcortical areas such as the thalamus, pallidum and putamen, and the visual cortex (superior occipital and lingual gyrus). Thalamic variability was associated with positive and general symptom scores, while variability in visual areas correlated with general symptoms such as poor impulse control and preoccupation (Table 3).

For autism patients, all regions showing significant changes had higher variability when compared to healthy controls, most significantly in the medial orbital and superior medial frontal gyrus, rectus and angular gyrus (Fig. 4B and Table 2). In particular, the variability of these default network regions was positively associated with the restricted, repetitive and stereotyped patterns of behaviour subscore (Table 3).

For ADHD patients, DMN regions involving the posterior cingulate cortex and angular gyrus all showed higher variability. In contrast, brain regions in the subcortical network, i.e. thalamus, showed lower variability in patients with ADHD compared to healthy control subjects (Fig. 4C and Table 2). Variability changes in posterior cingulate and frontal regions were associated with the severity of ADHD symptoms (Table 3).

Discussion

Brain regions with highest and lowest variability in healthy control subjects

We find that both primary sensory cortex and unimodal association cortex show very low variability because these regions are involved in unitary neural circuitry responsible for simple sensory functions. These regions are usually structurally connected more with regions belonging to the same modality (i.e. the same functional module) and their variability is small. In comparison, the transmodal areas (Mesulam, 1998), including the heteromodal association cortex and limbic regions, demonstrate high variability. These regions receive information from multiple sensory modalities and other heteromodal regions and are therefore responsible for more complex, integrated cognitive activities (Pearlson *et al.*, 1996; Bullard *et al.*, 2013). Consequently, these regions may participate in multiple functional communities at different times with resultant high temporal variability, or flexibility. Our results from resting state functional MRI are consistent with those obtained in task functional MRI by Bassett *et al.* (2013), who found that in a motor learning task primary sensorimotor and visual areas reconfigure little over time, while multimodal association regions reconfigure frequently.

Lastly, we note the relatively low variability of the DMN, including medial frontal gyrus and posterior cingulate/pre-cuneus, which is consistent with the strong functional connectivity within this network during resting state. These results are also in agreement with Power *et al.* (2011), who suggested that sensory-motor, visual and default mode systems are rather stationary.

Neural, electrophysiological and anatomical basis of variability

In terms of defining the neural basis of variability, Fig. 3 and Supplementary Table 4 show a negative correlation between the variability of a region and its BOLD activity/node degree. This indicated that the variability of a region is modulated by its BOLD activity. To facilitate information transmission with other regions, it is natural for a brain region to demonstrate greater BOLD activity to allow for high-level functional integration with other regions. Under these circumstances, this region will also have a high level of functional connectivity with other regions and, hence, higher degree. In this case, the temporal variability of the region of interest is expected to be low to maintain a high level of information transmission and functional integration. These results are consistent with previous studies (Bassett *et al.*, 2012; Zalesky *et al.*, 2012; Yu *et al.*, 2013), which showed that activity




Figure 4 Brain regions showing significant variability differences between patients with mental disorders and matched healthy controls. (A) Schizophrenia; (B) autism; (C) ADHD. Blue indicates that the variability of patients is lower than that of controls, and red indicates the opposite; see Table 2 for details.

and connectivity are related to one another in the resting state.

The negative correlation between the variability of a region of interest and the low frequency component of its BOLD signal (see Supplementary Fig. 5D, E and F and Supplementary Table 4) suggests that the low-frequency BOLD oscillation of a region facilitates information transmission and functional integration. This is essentially

because regions with more low-frequency components tend to synchronize more easily with other regions and thus have lower temporal variability. It is also consistent with the finding that low-frequency oscillations allow for integration of large neuronal networks (Buzsaki and Draguhn, 2004). In contrast, a region with more high-frequency components usually cannot synchronize effectively with other regions, thus demonstrating high variability.

Table 2 Significantly different regional variability

Brain region	Control	Patient	P-value	Brain region	Control	Patient	P-value
Schizophrenia				Vision regions: Vc < Vp			
DMN regions: Vc > Vp				(S) Lingual L	0.6451	0.6813	0.0018
Rectus L	0.7420	0.7036	0.0012	(S) Lingual R	0.6474	0.6776	0.0082
Hippocampus L	0.7458	0.7246	0.0103	(S) Occipital Sup L*	0.6421	0.6869	0.0003
Hippocampus R	0.7532	0.7210	0.0006	(S) Occipital Sup R	0.6463	0.6867	0.0011
(L) ParaHippocampal L*	0.7711	0.7304	0.00003	Subcortical regions: Vc < Vp			
(L) ParaHippocampal R*	0.7635	0.7114	3e-8	(S) Putamen R	0.6671	0.6967	0.0062
Parietal Inf L	0.7470	0.7221	0.0068	Pallidum R	0.6888	0.7186	0.0091
(L) Temporal Pole Mid L	0.7661	0.7318	0.0022	Thalamus R	0.7086	0.7405	0.0046
(L) Temporal Inf L*	0.7621	0.7263	0.0005	Other regions			
(L) Temporal Inf R	0.7779	0.7488	0.0020	Heschl L	0.6768	0.7111	0.0043
Autism				(L) Temporal Pole Sup L	0.7612	0.7357	0.0103
DMN regions: Vc < Vp				Supp Motor Area L	0.7488	0.7197	0.0048
(S) Frontal Sup Medial R	0.6878	0.7127	0.0018	Other regions: Vc < Vp			
(S) Frontal Med Orb L*	0.6873	0.7196	0.0002	Cingulum Ant L	0.7192	0.7404	0.0037
Frontal Med Orb R	0.7005	0.7274	0.0009	(L) Paracentral Lobule L	0.7909	0.8126	0.0036
Rectus L	0.7269	0.7531	0.0018	Subcortical regions: Vc > Vp			
Angular L	0.7205	0.7418	0.0029	Thalamus L	0.8002	0.7744	0.0038
ADHD				Thalamus R	0.8089	0.7835	0.0041
DMN regions: Vc < Vp				Other regions			
Angular L*	0.7479	0.7835	0.0002	(L) Frontal Sup Orb R	0.8203	0.7989	0.0026
Angular R	0.7445	0.7759	0.0030	(S) Insula L	0.6813	0.7084	0.0017
(S) Cingulum Post L*	0.7194	0.7591	0.0003	(S) Insula R	0.6671	0.6948	0.0035
Cingulum Post R*	0.7244	0.7636	0.0004	Supramarginal L	0.7596	0.7851	0.0023

Shown are data in patients with schizophrenia using the Taiwan (Dataset 1) and COBRE (Dataset 2) datasets; autism, using the NYU (Dataset 3) and UM (Dataset 4) datasets; and ADHD, using the PK dataset (Dataset 5) when compared with matched controls (related to Fig. 4). We used FDR ($q = 0.05$) for correction, and those regions which could survive Bonferroni correction ($P = 0.05$) are marked by an asterisk. The brain regions marked with 'L' or 'S' indicate that they are among the top 10 regions with the largest or smallest variability in matched healthy controls. Vp and Vc denotes mean variability for patient and control groups, respectively. Sup = superior; Orb = orbital; L = left; R = right; Post = posterior; Med = medial; Ant = anterior; Inf = inferior.

For the electrophysiological basis of variability, the variability of a region was modulated by the α band power of its EEG, as manifested by a positive correlation between variability of a region and its α power during the entire scan (Fig. 3A, C and Supplementary Table 5) across brain regions. This is consistent with the finding that α oscillation inhibits BOLD activity by showing a strong negative correlation with BOLD activity, especially in occipital, parietal and frontal cortices (Laufs et al., 2006; de Munck et al., 2007). In view of our finding that there is a negative correlation between BOLD activity and temporal variability (Fig. 3A and B), the observed positive correlation between α power and variability is therefore expected. Our results are based on correlation analysis across brain regions for each subject. In comparison, Chang et al. (2013) performed a correlation analysis across subjects for a given functional connectivity (i.e. between default-mode and dorsal attention networks). Further studies investigating

correlation between brain network variability and EEG signals at different bands (both across subjects and across brain regions) are needed to provide a full understanding of the electrophysiological basis of variability (Thompson et al., 2015).

In terms of the structural basis of variability, we found a negative correlation between variability and the ratio of intra- to inter-community structural connections (Fig. 3A and E). Thus, if a region is more structurally connected to nodes of the same functional community, then it will be more stably involved with this community and exhibit lower variability. In comparison, a region with structural connections to multiple communities would tend to frequently switch functional communities and demonstrate high variability.

As the resting state is a unconstrained condition that involves varying levels of mind-wandering, arousal, attention and vigilance (Chang and Glover, 2010), the

Table 3 Correlation between temporal variability that is significantly changed in patients

Brain region	Correlation coefficient	P-value	Score type
Schizophrenia			
Sum score			
Thalamus R	-0.2332	0.0129	Overall score
Thalamus R	-0.3347	0.0002	Positive score
Thalamus R	-0.2513	0.0073	General score
Lingual R	-0.3095	0.0009	Overall score
Lingual R	-0.3327	0.0003	General score
Temporal Pole Sup L	-0.2342	0.0103	Overall score
Temporal Pole Sup L	-0.2363	0.0087	General score
Subscore			
Thalamus R	-0.3037	0.0010	Unusual thought content (g)
Thalamus R	-0.3056	0.0007	Poor impulse control (g)
Lingual R	-0.2953	0.0014	Poor impulse control (g)
Lingual R	-0.2815	0.0023	Preoccupation (g)
Autism			
Frontal Med Orb L	0.2970	0.0163	ADI R RRB TOTAL C ^a
Rectus L	0.2739	0.0273	ADI R RRB TOTAL C ^a
Cingulum Ant L	0.3119	0.0114	ADI R RRB TOTAL C ^a
Angular L	0.3069	0.0098	ADOS STEREO BEHAV ^b
ADHD			
Frontal Sup Orb R	-0.2429	0.0177	Sum
Cingulum Post R	0.2598	0.011	Inattention
Frontal Sup Orb R	-0.3201	0.0016	Impulsive

^aADI R RRB TOTAL C: restricted, repetitive, and stereotyped patterns of behaviour subscore (C) total for autism diagnostic interview-revised.

^bADOS STEREO BEHAV: stereotyped behaviours and restricted interest total subscore of the classic Autism Diagnostic Observation Schedule (ADOS).

For schizophrenia, the results were obtained by integrating the Taiwan and COBRE datasets (FDR correction, $q = 0.05$) through meta-analysis. For autism, only the NYU dataset is used as the UM dataset has incomplete symptom scores. For ADHD, only PKU dataset is used as NYU dataset has incomplete scores. Sup = superior; Orb = orbital; L = left; R = right; Post = posterior; Med = medial; Ant = anterior; Inf = inferior.

temporal variability of functional brain networks derived from the BOLD functional MRI may be driven ultimately by changes in mental state. Further investigations are needed to explore how different levels of vigilance/mind-wandering can modulate BOLD/EEG activity and, in turn, the temporal variability of functional brain networks.

Disease-specific changes of variability

Resting state functional connectivity analysis has revealed changes in the intrinsic topographical organization of the brain in many psychiatric disorders, including schizophrenia, autism and ADHD (Castellanos *et al.*, 2008; Kennedy and Courchesne, 2008; Bassett *et al.*, 2012; Whitfield-Gabrieli and Ford, 2012). However, time-varying properties of brain networks in general mental disorders have been less investigated (Damaraju *et al.*, 2014; Schaefer *et al.*, 2014; Yu *et al.*, 2015). Our work has demonstrated disease-specific changes of variability in three mental disorders, which may provide clues to the dynamics underlying neuropathological profiles of different psychiatric

disorders and thereby contribute to the development of differential and diagnostic imaging.

Previous studies in schizophrenia have reported a hyper-activated and concomitantly hyper-connected DMN (Whitfield-Gabrieli *et al.*, 2009), which may mirror intensive self-reference and decreased attentional capacities in patients (Whitfield-Gabrieli *et al.*, 2009). In line with these findings, patients with schizophrenia exhibited decreased variability in those DMN regions associated with increased activity and connectivity in this disorder. Interaction between the DMN and task-positive networks is related to working memory and switching between an intrinsic and an extrinsic focus of attention (Weissman *et al.*, 2006; Whitfield-Gabrieli and Ford, 2012). Therefore, decreased DMN variability is associated with neurocognitive symptoms characteristic of schizophrenia, e.g. an exaggerated focus on one's own thoughts and a blurring of the boundary between internal and external worlds (Whitfield-Gabrieli *et al.*, 2009). Schizophrenia also demonstrates basic information processing deficits, particularly sensory gating (Bender *et al.*, 2007) associated with the thalamus. Increased variability in patients with schizophrenia in subcortical regions (such as thalamus,

Table 4 Disease-specific changes of variability for regions belonging to the DMN and subcortical areas for schizophrenia, autism and ADHD

	Schizophrenia	Autism	ADHD
DMN regions	Vp < Vc Hippocampus Parahippocampus Inferior parietal gyrus Inferior temporal gyrus Rectus	Vp > Vc Angular Rectus Medial frontal gyrus Superior medial frontal gyrus	Vp > Vc Posterior- cingulate Angular
Subcortical regions	Vp > Vc Thalamus Putamen		Vp < Vc Thalamus

'Vp' and 'Vc' denote variability of patients and healthy controls, respectively. Regions with significant changes are listed.

putamen, and pallidum; see Table 2) could therefore point to desynchronized basic filter modules associated with an inability to filter out irrelevant stimuli and a correspondingly diminished ability to focus attention (Freedman *et al.*, 1987). This is consistent with a recent study that reported hypo-connectivity of the putamen using a dynamic analysis (Damaraju *et al.*, 2014).

For autisms, a number of studies have found reduced DMN connectivity and activity (Kennedy and Courchesne, 2008) in resting state, which corresponds to the high variability of DMN regions we have observed. High variability in the DMN network suggests low functionality, i.e. disruption of resting state default-mode mediated cognition, which is mainly related to self-referential and theory of mind processing, inner speech, retrieving and manipulating memories, and future plans (Greicius *et al.*, 2003; Kennedy *et al.*, 2006; Garrity *et al.*, 2007). Therefore, self-referential thought may be reduced in autism (Cherkassky *et al.*, 2006) and directed more towards obsessive interests and sensory-environment processing than towards self-reflective activities (Crespi and Badcock, 2008).

Temporal variability analysis in ADHD patients revealed increased variability in regions of the DMN and concomitantly decreased variability in subcortical regions. Previous studies have also reported decreased DMN connectivity and integration (Castellanos *et al.*, 2008) in ADHD, as explained by the 'default-mode interference' hypothesis (Sonuga-Barke and Castellanos, 2007; Castellanos *et al.*, 2008). This theory suggests that the characteristic pattern of variability in performance in ADHD could be based on a dysfunctional synchronization in the DMN or its interactions with 'task-active' regions, e.g. decreased anti-correlations between the PCC and task-positive regions (Castellanos *et al.*, 2008). This is consistent with our finding of high variability in DMN regions such as the posterior cingulate in association with low functional connectivity, and may reflect default mode interference that contributes to attentional deficits in ADHD. In particular, the high variability associated with the posterior cingulate may be related its diminished volume (Carmona *et al.*,

2005) or decreased cortical thickness (Makris *et al.*, 2007) in ADHD, as well as in the precuneus.

Two interesting trends are worth noting. First, regions demonstrating extreme variability (either highest or lowest) in healthy controls are those most subject to change in mental disorders. As is shown in Table 2, half of the regions showing significant variability changes in the three disorders are among the top 10% of regions that have the highest or lowest variability in matched controls (11/19 in schizophrenia, 3/8 in autism, and 4/10 in ADHD). This indicates that regions at the two extremes of the axis of variability are unstable and tend to be affected by mental disorders. We find that regions with highest variability in controls (transmodal areas) always show a decrease in variability in disorders while regions with lowest variability in controls (primary sensory regions) tend to show an increase in disorders, suggesting abnormal functional integration tend to occur in primary sensory regions and transmodal areas in mental disorders.

Second, the three disorders studied showed a disease-specific and partly opposing pattern of altered variability in regions previously reported to be associated with symptoms. For example, schizophrenia and autism demonstrate opposing trends in variability changes in DMN regions compared with respective controls (Fig. 4 and Table 4), with posterior DMN regions affected in schizophrenia and anterior DMN regions affected in autism. This is consistent with the idea that schizophrenia and autism may represent contrasting pathologies (Crespi and Badcock, 2008). These disorders exhibit diametrically opposed phenotypes, or patterns of social brain development, such as social cognition, language, and behaviour, as well as local/global processing. Social cognition is thought to be underdeveloped in autistic-spectrum disorders, but hyperdeveloped in the psychotic spectrum (Crespi and Badcock, 2008). In addition, an opposing trend was also observed for variability changes in thalamus for schizophrenia and ADHD (Fig. 4 and Table 4).

Finally, although both autism and ADHD demonstrate changes in variability in DMN regions, the former involves mainly medial frontal areas, while the latter includes the

posterior cingulate. Autism spectrum disorders are characterized by deficits or inability to relate to other people and understand others' mental states (Assaf *et al.*, 2010). These theory of mind processes are supported by DMN activity, especially by the medial prefrontal cortex (Assaf *et al.*, 2010). The higher variability in medial frontal regions identified in autism indicates the low functionality of these regions, which may underlie the undermined theory of mind processes in autisms. In comparison, posterior cingulate dysfunction in ADHD is responsible for the disruption of the DMN that leads to attentional lapses (Leech and Sharp, 2014)

Relationship between variability and other measures and potential applications

We found that the temporal variability defined here is related to the concept of functional brain entropy proposed in our previous research (Yao *et al.*, 2013). Where a brain region exhibits large functional entropy this indicates a wider distribution of all its functional connections. In this case the region tends to maintain relatively high functional connectivity with many other regions (i.e. high network degree) and therefore demonstrates low temporal variability. Interestingly, regions showing a significant trend towards increased entropy with age, such as hippocampus/parahippocampal gyrus, olfactory cortex, and paracentral lobule, all have high temporal variability. On the other hand, regions demonstrating decreased entropy with age, such as the insular, have low temporal variability.

Bassett *et al.* (2011) have shown that flexibility of the brain is an important factor that predicts learning, and found correlations between the ability for the brain to learn and its flexibility (Bassett *et al.*, 2011, 2013, 2015). This suggests that the more often the brain switches connectivity patterns (i.e. more variable), the more flexible it is. Therefore, the temporal variability measure proposed in our paper may be a suitable indicator of the flexibility of a brain region, and could potentially be used to predict the outcome of learning. Our measure of brain variability may also have implications in rehabilitation, e.g. for patients who have had a stroke or glioma, or general brain injury. A brain region with more variability in global functional connectivity profile after brain injuries suggests its ability to participate in multiple communities, or neural systems, therefore this region is more likely to restore its function after rehabilitation or surgery. The variability measure proposed here consequently is also expected to reflect plasticity of the brain.

The influence of head movements

While it has been shown that head movements can influence estimates of functional connectivity (Satterthwaite *et al.*, 2013; Power *et al.*, 2014), evidence also indicates

that head movement only explains a small fraction of the variability in connectivity (Van Dijk *et al.*, 2012). Motion-associated differences in functional connectivity cannot be fully attributed to motion artefacts, but rather also reflect individual variability in functional organization (Zeng *et al.*, 2014). To evaluate the possible effect of head motion on temporal variability, we calculated the correlation between variability of a region and the head movements (Supplementary Table 6). Of the six functional MRI datasets (Datasets 1–6), only the patients in Dataset 4 (ADHD: NYU) show significant correlation between regional variability and head movement (nine regions survived correction; FDR, $q = 0.05$). We also listed regions with $P < 0.01$ (correlation between variability and head movement) for other datasets in Supplementary Table 6, and we found no consistent patterns across datasets. These results suggest that the effect of head movement is likely to be small and that the observed temporal variability in brain networks we have observed cannot be attributed to them, but may be driven by shifts in vigilance or mental state (Allen *et al.*, 2014). In the current study the NYU-dataset for the ADHD, which did show correlations with head movement was excluded from our analysis.

Acknowledgements

We would like to thank Prof. David Waxman, Prof. Junying Zhang and the reviewers for useful suggestions.

Funding

J.Z. is supported by the National Science Foundation of China (NSFC 61104143 and 61573107) and special Funds for Major State Basic Research Projects of China (2015CB856003). K.M.K. is supported by NSFC grant number 31530032. Z.L. is supported by NSFC 61201312 and 91530113, and Research Fund for the Doctoral Program of Higher Education of China (No. 2013 0203110017). X.L. is supported by the National Nature Science Foundation of China (No. 31571111). G.L. is supported by Key project of "12th Five-Year Plan" on medical science of PLA. J.F. is a Royal Society Wolfson Research Merit Award holder, partially supported by the National High Technology Research and Development Program of China (No. 2015AA020507) and the Key Project of Shanghai Science & Technology Innovation Plan (No. 15JC1400101), the National Centre for Mathematics and Interdisciplinary Sciences (NCMIS) of the Chinese Academy of Sciences, and the Key Program of the National Natural Science Foundation of China (No. 91230201).

Supplementary material

Supplementary material is available at *Brain* online.

References

- Allen EA, Damaraju E, Plis SM, Erhardt EB, Eichele T, Calhoun VD. Tracking whole-brain connectivity dynamics in the resting state. *Cereb Cortex* 2014; 24: 663–76.
- Assaf M, Jagannathan K, Calhoun VD, Miller L, Stevens MC, Sahl R, et al. Abnormal functional connectivity of default mode sub-networks in autism spectrum disorder patients. *Neuroimage* 2010; 53: 247–56.
- Baldassarre A, Lewis CM, Committeri G, Snyder AZ, Romani GL, Corbetta M. Individual variability in functional connectivity predicts performance of a perceptual task. *Proc Natl Acad Sci USA* 2012; 109: 3516–21.
- Bassett DS, Nelson BG, Mueller BA, Camchong J, Lim KO. Altered resting state complexity in schizophrenia. *Neuroimage* 2012; 59: 2196–207.
- Bassett DS, Wymbs NF, Porter MA, Mucha PJ, Carlson JM, Grafton ST. Dynamic reconfiguration of human brain networks during learning. *Proc Natl Acad Sci USA* 2011; 108: 7641–6.
- Bassett DS, Wymbs NF, Rombach MP, Porter MA, Mucha PJ, Grafton ST. Task-based core-periphery organization of human brain dynamics. *Plos Comput Biol* 2013; 9: e1003171.
- Bassett DS, Yang MZ, Wymbs NF, Grafton ST. Learning-induced autonomy of sensorimotor systems. *Nat Neurosci* 2015; 18: 744–51.
- Bender S, Weisbrod M, Resch F. Which perspectives can endophenotypes and biological markers offer in the early recognition of schizophrenia? *J Neural Transm* 2007; 114: 1199–215.
- Braun U, Schafer A, Walter H, Erk S, Romanczuk-Seiferth N, Haddad L, et al. Dynamic reconfiguration of frontal brain networks during executive cognition in humans. *Proc Natl Acad Sci USA* 2015; 112: 11678–83.
- Bullard SE, Griss M, Greene S, Gekker A. Encyclopedia of clinical neuropsychology. *Arch Clin Neuropsych* 2013; 28: 92.
- Buzsaki G, Draguhn A. Neuronal oscillations in cortical networks. *Science* 2004; 304: 1926–9.
- Calhoun VD, Miller R, Pearlson G, Adali T. The Chronnectome: time-varying connectivity networks as the next frontier in fMRI data discovery. *Neuron* 2014; 84: 262–74.
- Carmona S, Vilarroya O, Bielsa A, Tremols V, Soliva JC, Rovira M, et al. Global and regional gray matter reductions in ADHD: a voxel-based morphometric study. *Neurosci Lett* 2005; 389: 88–93.
- Castellanos FX, Margulies DS, Kelly C, Uddin LQ, Ghaffari M, Kirsch A, et al. Cingulate-precuneus interactions: a new locus of dysfunction in adult attention-deficit/hyperactivity disorder. *Biol Psychiat* 2008; 63: 332–7.
- Chang C, Glover GH. Time-frequency dynamics of resting-state brain connectivity measured with fMRI. *Neuroimage* 2010; 50: 81–98.
- Chang C, Liu ZM, Chen MC, Liu X, Duyn JH. EEG correlates of time-varying BOLD functional connectivity. *Neuroimage* 2013; 72: 227–36.
- Cheng W, Palaniyappan L, Li M, Kendrick KM, Zhang J, Luo Q, et al. Voxel-based, brain-wide association study of aberrant functional connectivity in schizophrenia implicates thalamocortical circuitry. *Npj Schizophrenia* 2015a; 1: 15016.
- Cheng W, Rolls ET, Gu HG, Zhang J, Feng JF. Autism: reduced connectivity between cortical areas involved in face expression, theory of mind, and the sense of self. *Brain* 2015b; 138: 1382–93.
- Cherkassky VL, Kana RK, Keller TA, Just MA. Functional connectivity in a baseline resting-state network in autism. *Neuroreport* 2006; 17: 1687–90.
- Crespi B, Badcock C. Psychosis and autism as diametrical disorders of the social brain. *Behav Brain Sci* 2008; 31: 241–61.
- Damaraju E, Allen EA, Belger A, Ford JM, McEwen S, Mathalon DH, et al. Dynamic functional connectivity analysis reveals transient states of dysconnectivity in schizophrenia. *Neuroimage Clin* 2014; 5: 298–308.
- de Munck JC, Goncalves SI, Huijboom L, Kuijer JPA, Pouwels PJW, Heethaar RM, et al. The hemodynamic response of the alpha rhythm: an EEG/fMRI study. *Neuroimage* 2007; 35: 1142–51.
- Deng W, Aimone JB, Gage FH. New neurons and new memories: how does adult hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* 2010; 11: 339–50.
- Fletcher ML, Chen WR. Neural correlates of olfactory learning: critical role of centrifugal neuromodulation. *Learn Mem* 2010; 17: 561–70.
- Freedman R, Adler LE, Gerhardt GA, Waldo M, Baker N, Rose GM, et al. Neurobiological studies of sensory gating in schizophrenia. *Schizophrenia Bull* 1987; 13: 669–78.
- Frost MA, Goebel R. Measuring structural-functional correspondence: spatial variability of specialised brain regions after macro-anatomical alignment. *Neuroimage* 2012; 59: 1369–81.
- Garrett DD, Kovacevic N, McIntosh AR, Grady CL. Blood oxygen level-dependent signal variability is more than just noise. *J Neurosci* 2010; 30: 4914–21.
- Garrett DD, Kovacevic N, McIntosh AR, Grady CL. The importance of being variable. *J Neurosci* 2011; 31: 4496–503.
- Garrity AG, Pearlson GD, McKiernan K, Lloyd D, Kiehl KA, Calhoun VD. Aberrant “default mode” functional connectivity in schizophrenia. *Am J Psychiat* 2007; 164: 450–7.
- Glahn DC, Laird AR, Ellison-Wright I, Thelen SM, Robinson JL, Lancaster JL, et al. Meta-analysis of gray matter anomalies in schizophrenia: application of anatomic likelihood estimation and network analysis. *Biol Psychiatry* 2008; 64: 774–81.
- Gould E, Reeves AJ, Graziano MS, Gross CG. Neurogenesis in the neocortex of adult primates. *Science* 1999; 286: 548–52.
- Greicius MD, Krasnow B, Reiss AL, Menon V. Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc Natl Acad Sci USA* 2003; 100: 253–8.
- Guo S, Kendrick KM, Yu R, Wang HLS, Feng J. Key functional circuitry altered in schizophrenia involves parietal regions associated with sense of self. *Hum Brain Mapp* 2014; 35: 123–39.
- Hutchison RM, Womelsdorf T, Allen EA, Bandettini PA, Calhoun VD, Corbetta M, et al. Dynamic functional connectivity: promise, issues, and interpretations. *Neuroimage* 2013a; 80: 360–78.
- Hutchison RM, Womelsdorf T, Gati JS, Everling S, Menon RS. Resting-state networks show dynamic functional connectivity in awake humans and anesthetized macaques. *Hum Brain Mapp* 2013b; 34: 2154–77.
- Jones DT, Vemuri P, Murphy MC, Gunter JL, Senjem ML, Machulda MM, et al. Non-stationarity in the “resting brain’s” modular architecture. *Plos One* 2012; 7: e3973.
- Kang J, Wang L, Yan CG, Wang JH, Liang X, He Y. Characterizing dynamic functional connectivity in the resting brain using variable parameter regression and Kalman filtering approaches. *Neuroimage* 2011; 56: 1222–34.
- Kawasaki K, Sheinberg DL. Learning to recognize visual objects with microstimulation in inferior temporal cortex. *J Neurophysiol* 2008; 100: 197–211.
- Kennedy DP, Courchesne E. The intrinsic functional organization of the brain is altered in autism. *Neuroimage* 2008; 39: 1877–85.
- Kennedy DP, Redcay E, Courchesne E. Failing to deactivate: resting functional abnormalities in autism. *Proc Natl Acad Sci USA* 2006; 103: 8275–80.
- Kopell NJ, Gritton HJ, Whittington MA, Kramer MA. Beyond the connectome: the dynamome. *Neuron* 2014; 83: 1319–28.
- Kucyi A, Davis KD. Dynamic functional connectivity of the default mode network tracks daydreaming. *Neuroimage* 2014; 100: 471–80.
- Laufs H, Holt JL, Elfont R, Krams M, Paul JS, Krakow K, et al. Where the BOLD signal goes when alpha EEG leaves. *Neuroimage* 2006; 31: 1408–18.
- Leech R, Sharp DJ. The role of the posterior cingulate cortex in cognition and disease. *Brain* 2014; 137: 12–32.

- Lippe S, Kovacevic N, McIntosh AR. Differential maturation of brain signal complexity in the human auditory and visual system. *Front Hum Neurosci* 2009; 3: 48.
- Liptak T. On the combination of independent tests. *Magyar Tud Akad Mat Kutato Int Kozl* 1958; 3: 171–97.
- Majeed W, Magnuson M, Hasenkamp W, Schwarb H, Schumacher EH, Barsalou L, et al. Spatiotemporal dynamics of low frequency BOLD fluctuations in rats and humans. *Neuroimage* 2011; 54: 1140–50.
- Makris N, Biederman J, Valera EM, Bush G, Kaiser J, Kennedy DN, et al. Cortical thinning of the attention and executive function networks in adults with Attention-Deficit/Hyperactivity disorder. *Cereb Cortex* 2007; 17: 1364–75.
- McIntosh AR, Kovacevic N, Itier RJ. Increased brain signal variability accompanies lower behavioral variability in development. *PLoS Comput Biol* 2008; 4: e1000106.
- Mesulam MM. From sensation to cognition. *Brain* 1998; 121: 1013–52.
- Misic B, Mills T, Taylor MJ, McIntosh AR. Brain noise is task dependent and region specific. *J Neurophysiol* 2010; 104: 2667–76.
- Mueller S, Wang DH, Fox MD, Yeo BTT, Sepulcre J, Sabuncu MR, et al. Individual variability in functional connectivity architecture of the human brain. *Neuron* 2013; 77: 586–95.
- Pearlson GD, Petty RG, Ross CA, Tien AY. Schizophrenia: a disease of heteromodal association cortex? *Neuropsychopharmacol* 1996; 14: 1–17.
- Power JD, Cohen AL, Nelson SM, Wig GS, Barnes KA, Church JA, et al. Functional network organization of the human brain. *Neuron* 2011; 72: 665–78.
- Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage* 2014; 84: 320–41.
- Protzner AB, Valiante TA, Kovacevic N, McCormick C, Mcandrews MP. Hippocampal signal complexity in mesial temporal lobe epilepsy: a noisy brain is a healthy brain. *Arch Ital Biol* 2010; 148: 289–97.
- Rademacher J, Burgel U, Geyer S, Schormann T, Schleicher A, Freund HJ, et al. Variability and asymmetry in the human precentral motor system - a cytoarchitectonic and myeloarchitectonic brain mapping study. *Brain* 2001; 124: 2232–58.
- Rakic P. Neurogenesis in adult primate neocortex: an evaluation of the evidence. *Nat Rev Neurosci* 2002; 3: 65–71.
- Samanez-Larkin GR, Kuhnen CM, Yoo DJ, Knutson B. Variability in nucleus accumbens activity mediates age-related suboptimal financial risk taking. *J Neurosci* 2010; 30: 1426–34.
- Satterthwaite TD, Elliott MA, Gerraty RT, Ruparel K, Loughead J, Calkins ME, et al. An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *Neuroimage* 2013; 64: 240–56.
- Schaefer A, Margulies DS, Lohmann G, Gorgolewski KJ, Smallwood J, Kiebel SJ, et al. Dynamic network participation of functional connectivity hubs assessed by resting-state fMRI. *Front Hum Neurosci* 2014; 8: 195.
- Schumann G, Loth E, Banaschewski T, Barbot A, Barker G, Buchel C, et al. The IMAGEN study: reinforcement-related behaviour in normal brain function and psychopathology. *Mol Psychiatr* 2010; 15: 1128–39.
- Seger CA, Cincotta CM. The roles of the caudate nucleus in human classification learning. *J Neurosci* 2005; 25: 2941–51.
- Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding subject-driven cognitive states with whole-brain connectivity patterns. *Cereb Cortex* 2012; 22: 158–65.
- Sonuga-Barke EJS, Castellanos FX. Spontaneous attentional fluctuations in impaired states and pathological conditions: a neurobiological hypothesis. *Neurosci Biobehav Rev* 2007; 31: 977–86.
- Sugiura M, Friston KJ, Willmes K, Shah NJ, Zilles K, Fink GR. Analysis of intersubject variability in activation: an application to the incidental episodic retrieval during recognition test. *Hum Brain Mapp* 2007; 28: 49–58.
- Tagliazucchi E, Laufs H. Decoding Wakefulness levels from typical fMRI resting-state data reveals reliable drifts between wakefulness and sleep. *Neuron* 2014; 82: 695–708.
- Thompson GJ, Pan WJ, Keilholz SD. Different dynamic resting state fMRI patterns are linked to different frequencies of neural activity. *J Neurophysiol* 2015; 114: 114–24.
- Van Dijk KRA, Sabuncu MR, Buckner RL. The influence of head motion on intrinsic functional connectivity MRI. *Neuroimage* 2012; 59: 431–8.
- Weissman DH, Roberts KC, Visscher KM, Woldorff MG. The neural bases of momentary lapses in attention. *Nat Neurosci* 2006; 9: 971–8.
- Whitfield-Gabrieli S, Ford JM. Default mode network activity and connectivity in psychopathology. *Ann Rev Clin Psychol* 2012; 8: 49–76.
- Whitfield-Gabrieli S, Thermenos HW, Milanovic S, Tsuang MT, Faraone SV, McCarley RW, et al. Hyperactivity and hyperconnectivity of the default network in schizophrenia and in first-degree relatives of persons with schizophrenia. *Proc Natl Acad Sci USA* 2009; 106: 1279–84.
- Williams ZM, Eskandar EN. Selective enhancement of associative learning by microstimulation of the anterior caudate. *Nat Neurosci* 2006; 9: 562–8.
- Yao Y, Lu WL, Xu B, Li CB, Lin CP, Waxman D, et al. The increase of the functional entropy of the human brain with age. *Sci Rep* 2013; 3: 2853.
- Yu QB, Erhardt EB, Sui J, Du YH, He H, Hjelm D, et al. Assessing dynamic brain graphs of time-varying connectivity in fMRI data: application to healthy controls and patients with schizophrenia. *Neuroimage* 2015; 107: 345–55.
- Yu QB, Sui J, Liu JY, Plis SM, Kiehl KA, Pearlson G, et al. Disrupted correlation between low frequency power and connectivity strength of resting state brain networks in schizophrenia. *Schizophr Res* 2013; 143: 165–71.
- Zalesky A, Fornito A, Cocchi L, Gollo LL, Breakspear M. Time-resolved resting-state brain networks. *Proc Natl Acad Sci USA* 2014; 111: 10341–6.
- Zalesky A, Fornito A, Egan GF, Pantelis C, Bullmore ET. The relationship between regional and inter-regional functional connectivity deficits in schizophrenia. *Hum Brain Mapp* 2012; 33: 2535–49.
- Zeng LL, Wang DH, Fox MD, Sabuncu M, Hu DW, Ge M, et al. Neurobiological basis of head motion in brain imaging. *Proc Natl Acad Sci USA* 2014; 111: 6058–62.